

## 1: Peri-Colace Uses, Side Effects & Warnings - [www.enganchecubano.com](http://www.enganchecubano.com)

*www.enganchecubano.com provides accurate and independent information on more than 24, prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.*

Docusate is a stool softener. It makes bowel movements softer and easier to pass. Senna is a laxative. It stimulates muscle movement in the intestines. The combination of docusate and senna is used to treat occasional constipation. Peri-Colace may also be used for other purposes not listed in this medication guide. Important Information Use this medication as directed on the label, or as your doctor has prescribed. Do not use the medication in larger amounts or for longer than recommended. You should not use this medication if you are allergic to Peri-Colace, or if you are also taking mineral oil. Do not take this medication for longer than 7 days in a row. Call your doctor if your constipation does not improve or if it gets worse. Stop taking this Peri-Colace and call your doctor at once if you have rectal bleeding , severe stomach pain, nausea and vomiting , or if you do not have a bowel movement. Do not use any other over-the-counter laxatives or other stool softener without first asking your doctor or pharmacist. Before taking this medicine You should not use this medication if you are allergic to Peri-Colace, or if you are also taking mineral oil. Ask a doctor or pharmacist about using Peri-Colace if you have: It is not known whether docusate and senna passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby. Ask a doctor before giving this medication to a child younger than 2 years old. How should I use Peri-Colace? Use this medication exactly as directed on the label, or as prescribed by your doctor. Do not use it in larger amounts or for longer than recommended. Take this medication with a full glass of water. It may be best to take this medication at night or at bedtime. Peri-Colace should cause you to have a bowel movement within 6 to 12 hours. Do not take this medication for longer than 7 days in a row, unless your doctor tells you to. Call your doctor if your constipation does not improve or if it gets worse after taking Peri-Colace. Store Peri-Colace at room temperature away from moisture and heat. What happens if I miss a dose? Since Peri-Colace is taken as needed, you are not likely to be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take extra medicine to make up the missed dose. What happens if I overdose? Seek emergency medical attention if you think you have used too much of Peri-Colace. Overdose symptoms may include nausea, vomiting, stomach pain, or diarrhea. What should I avoid while using Peri-Colace? Docusate or senna may be contained in other medicines available over the counter. If you take certain products together you may accidentally take too much of a certain medicine. Read the label of any other medicine you are using to see if it contains docusate or senna. Peri-Colace side effects Get emergency medical help if you have any of these signs of an allergic reaction: Stop using Peri-Colace and call your doctor at once if you have a serious side effect such as:

**2: "Docosate sodium with senna" Keyword Found Websites Listing | Keyword Suggestions**

*How to use Docosate SODIUM-SENNA. The effects of some drugs can change if you take other drugs or herbal products at the same time. Senna with Docosate Sodium mg mg tablet. Senna.*

Other names Why is this medication prescribed? Senna is used on a short-term basis to treat constipation. It also is used to empty the bowels before surgery and certain medical procedures. Senna is in a class of medications called stimulant laxatives. It works by increasing activity of the intestines to cause a bowel movement. How should this medicine be used? Senna comes as a liquid, powder, granules, chewable pieces, and tablets to take by mouth. It may be taken once or twice daily. Senna normally causes a bowel movement within 6 to 12 hours, so it may be taken at bedtime to produce a bowel movement the next day. Do not take senna for more than 1 week without talking to your doctor. Follow the directions on your package or prescription label carefully, and ask your doctor or pharmacist to explain any part you do not understand. Take senna exactly as directed. Frequent or continued use of senna may make you dependent on laxatives and cause your bowels to lose their normal activity. If you do not have a regular bowel movement after taking senna, do not take any more medication and talk to your doctor. Other uses for this medicine This medication may be prescribed for other uses; ask your doctor or pharmacist for more information. What special precautions should I follow? Before taking senna, tell your doctor and pharmacist if you are allergic to senna, any other medications, or any of the ingredients in these senna products. Ask your pharmacist for a list of the ingredients. Be sure to mention: Your doctor may need to change the doses of your medications or monitor you carefully for side effects. If you become pregnant while taking senna, call your doctor. Older adults should not usually take senna products over a long period of time because they are not as safe as other medications that can be used to treat the same condition. What special dietary instructions should I follow? A regular diet and exercise program is important for regular bowel function. Eat a high-fiber diet and drink plenty of liquids eight glasses each day as recommended by your doctor. What should I do if I forget a dose? This medication usually is taken as needed. If your doctor has told you to take senna regularly, take the missed dose as soon as you remember it. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. Do not take a double dose to make up for a missed one. What side effects can this medication cause? Senna may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away:

### 3: Senna: MedlinePlus Drug Information

*Peri-Colace may also be used for other purposes not listed in this medication guide. Important Information Use this medication as directed on the label, or as your doctor has prescribed.*

**Descriptions** Oral laxatives are medicines taken by mouth to encourage bowel movements to relieve constipation. There are several different types of oral laxatives and they work in different ways. Since directions for use are different for each type, it is important to know which one you are taking. The different types of oral laxatives include:

- Bulk-formers**—Bulk-forming laxatives are not digested but absorb liquid in the intestines and swell to form a soft, bulky stool. The bowel is then stimulated normally by the presence of the bulky mass. Some bulk-forming laxatives, like psyllium and polycarbophil, may be prescribed by your doctor to treat diarrhea.
- Hyperosmotics**—Hyperosmotic laxatives encourage bowel movements by drawing water into the bowel from surrounding body tissues. This provides a soft stool mass and increased bowel action. There are three types of hyperosmotic laxatives taken by mouth—the saline, the lactulose, and the polymer types: The saline type is often called "salts. They are not used for long-term or repeated correction of constipation. With smaller doses than those used for the laxative effect, some saline laxatives are used as antacids. The information that follows applies only to their use as laxatives. Sodium phosphate may also be prescribed for other conditions as determined by your doctor. The lactulose type is a special sugar-like laxative that works the same way as the saline type. However, it produces results much more slowly and is often used for long-term treatment of chronic constipation. Lactulose may sometimes be used in the treatment of certain medical conditions to reduce the amount of ammonia in the blood. The polymer type is a polyglycol polyethylene glycol, a large molecule that causes water to be retained in the stool; this will soften the stool and increase the number of bowel movements. It is used for short periods of time to treat constipation.
- Lubricants**—Lubricant laxatives, such as mineral oil, taken by mouth encourage bowel movements by coating the bowel and the stool mass with a waterproof film. This keeps moisture in the stool. The stool remains soft and its passage is made easier.
- Stimulants**—Stimulant laxatives, also known as contact laxatives, encourage bowel movements by acting on the intestinal wall. They increase the muscle contractions that move along the stool mass. Stimulant laxatives are a popular type of laxative for self-treatment. However, they also are more likely to cause side effects. One of the stimulant laxatives, dehydrocholic acid, may also be used for treating certain conditions of the biliary tract.
- Stool softeners emollients**—Stool softeners encourage bowel movements by helping liquids mix into the stool and prevent dry, hard stool masses. This type of laxative has been said not to cause a bowel movement but instead allows the patient to have a bowel movement without straining.

**Combinations**—There are many products that you can buy for constipation that contain more than one type of laxative. For example, a product may contain both a stool softener and a stimulant laxative. In general, combination products may be more likely to cause side effects because of the multiple ingredients. In addition, they may not offer any advantage over products containing only one type of laxative. If you are taking a combination laxative, make certain you know the proper use and precautions for each of the different ingredients. Most laxatives except saline laxatives may be used to provide relief:

**4: L.R. by Cristina Sanchez on Prezi**

*Davis's Drug Guide App + Web powered by Unbound Medicine provides up-to-date information on + trade name and generic drugs. View on the web or download App to your iPhone, iPad, or Android smartphone or tablet.*

Docusate is a stool softener. It makes bowel movements softer and easier to pass. Senna is a laxative. It stimulates muscle movement in the intestines. The combination of docusate and senna is used to treat occasional constipation. Docusate and senna may also be used for other purposes not listed in this medication guide. Use this medication as directed on the label, or as your doctor has prescribed. Do not use the medication in larger amounts or for longer than recommended. You should not use this medication if you are allergic to docusate and senna, or if you are also taking mineral oil. Do not take this medication for longer than 7 days in a row. Call your doctor if your constipation does not improve or if it gets worse. Stop taking this docusate and senna and call your doctor at once if you have rectal bleeding, severe stomach pain, nausea and vomiting, or if you do not have a bowel movement. Do not use any other over-the-counter laxatives or other stool softener without first asking your doctor or pharmacist. Ask a doctor or pharmacist about using docusate and senna if you have: It is not known whether docusate and senna passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby. Ask a doctor before giving this medication to a child younger than 2 years old. [Back to Top Docusate-Senna Side Effects](#) Get emergency medical help if you have any of these signs of an allergic reaction: Stop using docusate and senna and call your doctor at once if you have a serious side effect such as: Less serious side effects may include: This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect. [Back to Top Docusate-Senna Interactions](#) Do not use any other over-the-counter laxatives or other stool softener without first asking your doctor or pharmacist. Docusate or senna may be contained in other medicines available over the counter. If you take certain products together you may accidentally take too much of a certain medicine. Read the label of any other medicine you are using to see if it contains docusate or senna. There may be other drugs that can interact with docusate and senna. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor. [Back to Top Docusate-Senna Dosage](#) Use this medication exactly as directed on the label, or as prescribed by your doctor. Do not use it in larger amounts or for longer than recommended. Take this medication with a full glass of water. It may be best to take this medication at night or at bedtime. Docusate and senna should cause you to have a bowel movement within 6 to 12 hours. Do not take this medication for longer than 7 days in a row, unless your doctor tells you to. Call your doctor if your constipation does not improve or if it gets worse after taking docusate and senna. Store docusate and senna at room temperature away from moisture and heat. Seek emergency medical attention if you think you have used too much of this medicine. Overdose symptoms may include nausea, vomiting, stomach pain, or diarrhea. Since docusate and senna is taken as needed, you are not likely to be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take extra medicine to make up the missed dose. Copyright Cerner Multum, Inc.

**5: Docusate and Senna: Indications, Side Effects, Warnings - [www.enganchecubano.com](http://www.enganchecubano.com)**

*The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects, nor should it be construed to indicate that use of a particular drug is.*

Uses of Docusate and Senna: It is used to treat hard stools constipation. If you have an allergy to docusate, senna, or any other part of docusate and senna. If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives ; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs. If you have any of these health problems: Bowel block, belly pain, upset stomach , rectal bleeding , throwing up, or change in bowel habits lasting longer than 2 weeks. If you are taking mineral oil. This is not a list of all drugs or health problems that interact with docusate and senna. Tell your doctor and pharmacist about all of your drugs prescription or OTC, natural products, vitamins and health problems. You must check to make sure that it is safe for you to take docusate and senna with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor. What are some things I need to know or do while I take Docusate and Senna? Tell all of your health care providers that you take docusate and senna. This includes your doctors, nurses, pharmacists, and dentists. Do not use docusate and senna for more than 1 week unless told to do so by your doctor. Do not use other laxatives or stool softeners unless told to do so by the doctor. If you have rectal bleeding or you do not have a bowel movement after using docusate and senna, talk with your doctor. Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using docusate and senna while you are pregnant. Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby. How is this medicine Docusate and Senna best taken? Use docusate and senna as ordered by your doctor. Read all information given to you. Follow all instructions closely. Follow how to take docusate and senna as you have been told by your doctor. Do not use more than you were told to use. Take at bedtime if taking once a day. Take with a full glass of water. Do not take other drugs within 2 hours of docusate and senna. What do I do if I miss a dose? If you take docusate and senna on a regular basis, take a missed dose as soon as you think about it. If it is close to the time for your next dose, skip the missed dose and go back to your normal time. Do not take 2 doses at the same time or extra doses. Many times docusate and senna is taken on an as needed basis. Do not take more often than told by the doctor. Dosage Information in more detail What are some side effects that I need to call my doctor about right away? Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. What are some other side effects of Docusate and Senna? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if you have any side effects that bother you or do not go away. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may also report side effects at [http:](http://) If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened. Store at room temperature. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. Consumer information use If your symptoms or health problems do not get better or if they become worse, call your doctor. Keep a list of all your drugs prescription, natural products, vitamins , OTC with you. Give this list to your doctor. Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions

about docusate and senna, please talk with your doctor, nurse, pharmacist, or other health care provider. Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

## 6: "Senna and docusate" Keyword Found Websites Listing | Keyword Suggestions

*senna (sennosides; Senokot, Senokot EXTRA and others) is a natural medicine used to treat constipation. Side effects, drug interactions, dosage, and pregnancy safety information should be reviewed prior to taking this supplement.*

## 7: SENNA (SENNOSIDES)

*Docusate sodium/senna | Davis's Drug Guide [www.enganchecubano.com](http://www.enganchecubano.com) docusate sodium/senna is a topic covered in the Davis's Drug Guide. To view the entire topic, please sign in or purchase a subscription.*

## 8: Docusate-Senna - Side Effects, Dosage, Interactions - Drugs - Everyday Health

*Docusate / Senna Dosage Guide with Precautions - [www.enganchecubano.com](http://www.enganchecubano.com) [www.enganchecubano.com](http://www.enganchecubano.com) Docusate-senna should generally not be used for the treatment of chronic constipation or a period greater than one week, unless otherwise directed by a physician.*

*Of farms and fish: 1910-1915 Wizard world history by ankur sharma Great Works of Art and What Makes Them Great Irish thoroughbred nora roberts Photoelectrochemistry and Photovoltaics of Layered Semiconductors (Physics and Chemistry of Materials wit Physical Examinations and Health Assessment, Third Edition (Student Laboratory Manual) Classification of remedies The Politics of Ancient Israel (Library of Ancient Israel) Blakes bike and other synenergezetic sculpture Play like a girl by Kristine Kathryn Rusch Future of Central America Airplanes (Worlds Fastest Machines) May you be in heaven before the devil knows youre dead Ralph Compton Trail to Cottonwood Falls (Ralph Compton Western Series) 19-4. U-tube manometer, open type 332 Trees, woods and forests Chalabi under seige, May 2003 11 Flash for wildlife photography in the field 58 Gravitation and Modern Cosmology (Ettore Majorana International Science Series: Physical Sciences) Fancy Nancy Loves! Loves! Loves! (Fancy Nancy) The Warsaw diary of Chaim A. Kaplan. Vampire the masquerade revised edition Painting on glass Learn indian stock market basics U.s history the americans ch29 sect1 Adult learning in America Peter and the Wolf, from Serge Prokoffef IV. How Friar Giles praised obedience more than prayer 226 The broomstick train. Better food for pregnancy Birds of the Solomons, Vanuatu New Caledonia The same answer everywhere, that they might be received among the faithful, but not again exercise their Stopping mass killings in Africa Old Paths And Legends Of New England Engineering graphics and design grade 12 workbook memo Bmw x3 2007 manual Dell powervault md3000i manual Saving angel jl weil A Fortune branches out Assistance to veterans of Vietnam or elsewhere in obtaining suitable employment.*